



Ring Ouzel. Winter. Adult. Male (29-XII)

### RING OUZEL (Turdus torquatus)

### **IDENTIFICATION**

24-27 cm. **Adult male** black in breeding plumage; pectoral band pure white; yellow bill with dark tip; in non breeding plumage in **winter** with greyish edges on feathers and dark bill with yellowish tip. **Female** with dark grey plumage; pectoral band dark brown. Some 1st year females without pale pectoral band.



### SIMILAR SPECIES

The white pectoral band makes this species unmistakable. **Females** without white band recall a **Spotless Starling** with has shorter tail and thinner bill; **Female Blue Rock Thrush** has similar size, but different pattern on flanks and underparts; **Blackbird** has wing feathers without pale edges.









#### **SEXING**

In adult birds, breeding male with a white or light brownish pectoral band; head and neck black or brownish-black; in autumn feathers on head and neck with brownish edges; body feathers with a narrow pale edge (about 1 mm). Breeding female with obscure brownish-white pectoral band; head and neck dak brownish; in autumn feathers on head and neck with paler edges; body feathers with a wide brownish edge (about 2 mm) (CAUTION: some adult females are very similar to males). In 1st year autumn/2nd year spring birds, male with an evident pectoral band, paler than underparts; female with poor marked pectoral band, with the same colour than underparts. Juveniles cannot be sexed.





Ring Ouzel. Adult. Sexing. Pattern of band on breast: left male; right female.





Ring Ouzel. 1st year. Sexing. Pattern of band on breast: left male; right female.

#### **AGEING**

3 types of age can be recognized:

**Juvenile** has body feathers with pale tips; pale chin and throat; pale mottled on tips of lesser and median coverts; greater coverts with large whitish edge.

**1st year autumn/2ne year spring** with moult limit between the inner changed greater coverts (narrow pale edge) and the outermost unmoulted; tail feathers with pointed tips; some **females** with unmarked pectoral band.


**Adult** with no moult limits on wing; all the greater coverts with narrow edge; tail feathers with rounded tips.





Ring Ouzel. Ageing. Pattern of breast: left adult female; right juvenile.






Ring Ouzel. Ageing. Pattern of wing coverts: top adult female; bottom juvenile.








Ring Ouzel. Female. Ageing. Pattern of band on breast: left adult; right 1st year.





Ring Ouzel. Male. Ageing. Moult limit on wing coverts: top adult; bottom 1st year.





Ring Ouzel. Ageing. Pattern of tips of tail feathers: left adult; right 1st year.






Ring Ouzel. Female. Ageing. Moult limit on wing coverts: top adult; bottom 1st year.

#### **MOULT**

Complete **postbreeding** moult, usually finished in September before migration. Partial **postjuvenile** moult including body feathers, lesser and median coverts and a variable number of greater coverts; some birds can moult tetials; usually finished in October.



R i n g O u z e l . Extent of postjuvenile moult.



## **PHENOLOGY**



## **STATUS IN ARAGON**

Breeding only in the Pyrenees but wintering in lower areas of the Region, mainly in *Juniperus thurifera* forests of the Ebro Valley and Teruel.





Ring Ouzel. Winter. Adult. Head pattern: top male (29-XII); bottom female (10-I)




Ring Ouzel. Winter. Adult. Female (10-I)




Ring Ouzel.
Winter. 1st
year. Head
pattern: top
male (01XI); bottom
female ()



Ring Ouzel. Winter. 1st year. Male (01-XI)





Ring Ouzel. Winter. 1st year. Female ()


Ring Ouzel. Winter. Adult. Crown pattern: left male (29-XII); right female (10-I)







Ring Ouzel. Winter1st year. Crown pattern: left male (01-XI); right female ()







**NO ÌMAGE** 

Ring Ouzel. Winter. Adult. Breast pattern: left male (29-XII); right female (10-I)




Ring Ouzel. Winter1st year. Breast pattern: left male (01-XI); right female ()



Ring Ouzel. Winter. Adult. Neck pattern: left male (29-XII); right female (10-I)





Ring Ouzel. Winter1st year. Neck pattern: left male (01-XI); right female ()





Ring Ouzel. Winter. Adult. Upperparts pattern: left male (29-XII); right female (10-I)








Ring Ouzel. Winter1st year. Upperparts pattern: left male (01-XI); right female ()





Ring Ouzel. Winter. Adult. Tail pattern: left male (29-XII); right female (10-I)





Ring Ouzel. Winter1st year. Tail pattern: left male (01-XI); right female ()




Ring Ouzel. Winter. Adult. Male: pattern of wing (29-XII).



Ring Ouzel. Winter. Adult. Female: pattern of wing (10-I).



Ring Ouzel. Winter. 1st year. Male: pattern of wing (01-XI)



Ring Ouzel. Winter. 1st year. Female: pattern of wing ()